

R-evolution: Improving perturbative QCD

André H. Hoang,¹ Ambar Jain,² Ignazio Scimemi,³ and Iain W. Stewart²

¹*Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), Föhringer Ring 6, 80805 München, Germany*

²*Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139*

³*Departamento de Física Teórica II, Universidad Complutense de Madrid, 28040 Madrid, Spain*

Perturbative QCD results in the $\overline{\text{MS}}$ scheme can be dramatically improved by switching to a scheme that accounts for the dominant power law dependence on the factorization scale in the operator product expansion. We introduce the “MSR scheme” which achieves this in a Lorentz and gauge invariant way. The MSR scheme has a very simple relation to $\overline{\text{MS}}$, and can be easily used to reanalyze $\overline{\text{MS}}$ results. Results in MSR depend on a cutoff parameter R , in addition to the μ of $\overline{\text{MS}}$. R variations can be used to independently estimate i) the size of power corrections, and ii) higher order perturbative corrections (much like μ in $\overline{\text{MS}}$). We give two examples at three-loop order, the ratio of mass splittings in the B^*-B and D^*-D systems, and the Ellis-Jaffe sum rule as a function of momentum transfer Q in deep inelastic scattering. Comparing to data, the perturbative MSR results work well even for $Q \sim 1$ GeV, and power corrections are reduced compared to $\overline{\text{MS}}$.

Introduction and Formalism

The operator product expansion (OPE) is an important tool for QCD. In hard scattering processes two important scales are Q , a large moment transfer or mass, and Λ_{QCD} , the scale of nonperturbative matrix elements. The Wilsonian OPE introduces a factorization scale Λ^f , where $\Lambda_{\text{QCD}} < \Lambda^f < Q$, and expands in Λ_{QCD}/Q . Consider a dimensionless observable σ whose OPE is

$$\sigma = C_0^W(Q, \Lambda^f) \theta_0^W(\Lambda^f) + C_1^W(Q, \Lambda^f) \frac{\theta_1^W(\Lambda^f)}{Q^p} + \dots \quad (1)$$

The $C_{0,1}^W$ are dimensionless Wilson coefficients containing contributions from momenta $k > \Lambda^f$ with perturbative expansions in α_s , and $\theta_{0,1}^W = \langle \mathcal{O}_{0,1} \rangle_W$ are nonperturbative matrix element with mass dimensions 0 and p , containing contributions from $k < \Lambda^f$. If $C_{0,1}^W(Q, \Lambda^f)$ are expanded they contain an infinite series of terms, $(\Lambda^f/Q)^n$, modulo $\ln^m(\Lambda^f/Q)$ terms, and this reflects the fact that $C_{0,1}^W$ only include contributions from momenta $k > \Lambda^f$. The Wilsonian OPE provides a clean separation of momentum scales, but can be technically challenging to implement. In particular, it is difficult to define Λ^f and retain gauge symmetry and Lorentz invariance, and perturbative computations beyond one-loop are atrocious.

A popular alternative is the OPE with dimensional regularization and the $\overline{\text{MS}}$ scheme, which preserves the symmetries of QCD and provides powerful techniques for multiloop computations. In this case Eq. (1) becomes

$$\sigma = \bar{C}_0(Q, \mu) \bar{\theta}_0(\mu) + \bar{C}_1(Q, \mu) \frac{\bar{\theta}_1(\mu)}{Q^p} + \dots, \quad (2)$$

where μ is the renormalization scale and bars are used for $\overline{\text{MS}}$ quantities. In $\overline{\text{MS}}$ the $\bar{C}_i$ are simple series in α_s ,

$$\bar{C}_i(Q, \mu) = 1 + \sum_{n=1}^{\infty} b_n \left(\frac{\mu}{Q} \right) \left[\frac{\alpha_s(\mu)}{(4\pi)} \right]^n, \quad (3)$$

with coefficients $b_n(\mu/Q) = \sum_{k=0} b_{nk} \ln^k(\mu/Q)$ containing only $\ln \mu/Q$. We will always rescale σ and the matrix

elements $\bar{\theta}_i$ such that $\bar{C}_i = 1$ at tree level. In $\overline{\text{MS}}$ all power law dependence on Q is manifest and unique in each term of Eq. (2). Also simple renormalization group equations in μ , like $d \ln \bar{C}_0(Q, \mu)/d \ln \mu = \bar{\gamma}[\alpha_s(\mu)]$, can be used to sum large logs in Eq. (2) if $Q \gg \Lambda_{\text{QCD}}$.

$C_i^W(Q, \Lambda^f)$ and $\bar{C}_i(Q, \mu)$ are related to each other in perturbation theory, so Eqs. (1) and (2) are just the same OPE in two different schemes. The renormalization scale μ in $\overline{\text{MS}}$ plays the role of Λ^f . This is precisely true for logarithmic contributions, $\ln \mu \leftrightarrow \ln \Lambda^f$, and here the Wilsonian picture of scale separation in $\bar{C}_i$ and $\bar{\theta}_i$ carries over. However, the same is not true for power law dependences on Λ^f . $\overline{\text{MS}}$ integrations are carried out over all momenta, so the $\bar{C}_i$ actually contain some contributions from arbitrary small momenta, and the $\bar{\theta}_i$ have contributions from arbitrary large momenta. For the power law terms there is no explicit scale separation in $\overline{\text{MS}}$, and correspondingly no powers of μ appear in Eq. (3). While this simplifies higher order computations, it is known to lead to factorial growth in the perturbative coefficients. For $\bar{C}_0$, one has $b_{n+1}(\mu/Q) \simeq (\mu/Q)^p n! [2\beta_0/p]^n Z$ at large n [1], for constant Z . In practice this sometimes leads to poor convergence already at one or two loop order in QCD. This poor behavior is canceled by corresponding instabilities in $\bar{\theta}_1$, and is referred to as an order- p infrared renormalon in $\bar{C}_0$ canceling against an ultraviolet renormalon in $\bar{\theta}_1$ [2, 3, 4]. The cancellation reflects the fact that the $\overline{\text{MS}}$ OPE does not strictly separate momentum scales.

The OPE can be converted to a scheme that removes this poor behavior, but still retains the simple computational features of $\overline{\text{MS}}$. Consider defining a new “ R -scheme” for C_0 by subtracting a perturbative series

$$C_0(Q, R, \mu) = \bar{C}_0(Q, \mu) - \delta C_0(Q, R, \mu),$$

$$\delta C_0(Q, R, \mu) = \left(\frac{R}{Q} \right)^p \sum_{n=1}^{\infty} d_n \left(\frac{\mu}{R} \right) \left[\frac{\alpha_s(\mu)}{(4\pi)} \right]^n, \quad (4)$$

with $d_n(\mu/R) = \sum_{k=0} d_{nk} \ln^k(\mu/R)$. If for large n the coefficients d_n are chosen to have the same behavior as b_n ,

so $d_{n+1}(\mu/Q) \simeq (\mu/R)^p n! [2\beta_0/p]^n Z$, then the factorial growth in $\bar{C}_0(Q, \mu)$ and $\delta C_0(Q, R, \mu)$ cancel,

$$C_0(Q, R, \mu) \sim \left[\frac{\mu^p}{Q^p} - \frac{R^p}{Q^p} \frac{\mu^p}{R^p} \right] \sum_n n! \left[\frac{2\beta_0}{p} \right]^n Z. \quad (5)$$

Thus the R -scheme introduces power law dependence on the cutoff, $(R/Q)^p$, in $C_0(Q, R, \mu)$, which captures the dominant $(\Lambda^f/Q)^p$ behavior of the Wilsonian C_0^W . In practice this improves the convergence in C_0 even at low orders in the α_s series. The dominant effect of this change is compensated by a scheme change to $\bar{\theta}_1$, $\bar{\theta}_1(\mu) = \theta_1(R, \mu) - [Q^p \delta C_0(Q, R, \mu)] \bar{\theta}_0(\mu)$, and the new θ_1 will exhibit improved stability. In the R -scheme the OPE becomes

$$\sigma = C_0(Q, R, \mu) \bar{\theta}_0(\mu) + \bar{C}_1(Q, \mu) \theta_1(R, \mu)/Q^p + \bar{C}'_1(Q, \mu) \theta'_1(R, \mu)/Q^p + \dots, \quad (6)$$

where $\theta'_1 = [Q^p \delta C_0] \bar{\theta}_0$ and $\bar{C}'_1 = 1 - \bar{C}_1 \sim \alpha_s$. Both C_0 and θ_1 are free of order- p renormalons. The series in $\bar{C}'_1 \theta'_1$ is Borel summable. In all examples below $\bar{\theta}_0$ is also renormalon free. The above procedure may be repeated for higher renormalons and the higher power terms in the OPE indicated by ellipses, to improve the behavior of these terms as well. At the order at which we work, we will consistently set $\bar{C}_1 = 1$ and drop θ'_1 in the following.

To setup an appropriate R -scheme it remains to define the d_n . In the renormalon literature such scheme changes are well known for masses. For OPE predictions a “renormalon subtraction” (RS) scheme has been implemented in Ref. [5]. In RS an approximate result for the residue of the leading Borel renormalon pole is used to define the d_n , which adds a source of uncertainty.

For our analysis we define the “MSR” scheme for C_0 by simply taking the coefficients of the subtraction to be exactly the $\overline{\text{MS}}$ coefficients. In general it is more convenient to use $\ln \bar{C}_0$ rather than $\bar{C}_0$, since this simplifies renormalization group equations. Writing the series as

$$\ln \bar{C}_0(Q, \mu) = \sum_{n=1}^{\infty} a_n(\mu/Q) \left[\frac{\alpha_s(\mu)}{(4\pi)} \right]^n, \quad (7)$$

with $a_n(\mu/Q) = \sum_{k=0} a_{nk} \ln^k \mu/Q$ we define the MSR scheme by the series

$$\ln C_0(Q, R, \mu) \equiv \sum_{n=1}^{\infty} \left\{ a_n \left(\frac{\mu}{Q} \right) - \frac{R^p}{Q^p} a_n \left(\frac{\mu}{R} \right) \right\} \frac{\alpha_s^n(\mu)}{(4\pi)^n}. \quad (8)$$

This definition still cancels the order- p renormalon for large n , as in Eq. (5). It yields the very simple relation

$$C_0(Q, R, \mu) = \bar{C}_0(Q, \mu) [\bar{C}_0(R, \mu)]^{-(R/Q)^p}, \quad (9)$$

which must be expanded order-by-order in $\alpha_s(\mu)$ to remove the renormalon. Thus the coefficient $C_0(Q, R, \mu)$ for the MSR scheme is obtained directly from the $\overline{\text{MS}}$ result. Note $C_0(Q, Q, \mu) = 1$ to all orders. The appropriate p is obtained from the $\overline{\text{MS}}$ OPE by $p = \text{dimension}(\bar{\theta}_1) -$

dimension($\bar{\theta}_0$). MSR preserves gauge invariance, Lorentz symmetry, and the simplicity of $\overline{\text{MS}}$.

The appropriate values for R in Eqs. (4,6,9) are constrained by power counting and the structure of large logs in the OPE. The power counting $\bar{\theta}_1 \sim \Lambda_{\text{QCD}}^p$ implies $\theta_1 \sim \Lambda_{\text{QCD}}^p$, so for the matrix element we need $R \sim \mu \gtrsim \Lambda_{\text{QCD}}$ (meaning a larger value where perturbation theory for the OPE still converges), which minimizes $\ln(\mu/\Lambda_{\text{QCD}})$ and $\ln(\mu/R)$ terms in $\theta_1(R, \mu, \Lambda_{\text{QCD}})$. On the other hand, $C_0(Q, R, \mu)$ has $\ln(\mu/Q)$ and $\ln(\mu/R)$ terms, and for $R \sim \Lambda_{\text{QCD}}$ no choice of μ avoids large logs. For $R \sim \mu \sim Q$ we can minimize the logs in $C_0(Q, R, \mu)$, but not in $\theta_1(R, \mu, \Lambda_{\text{QCD}})$. When the OPE is carried out in $\overline{\text{MS}}$ this problem is dealt with using a μ -RGE to sum large logs between Q and Λ_{QCD} . For MSR we must use R -evolution, an RGE in the R variable [6]. The appropriate R -RGE is formulated with $\mu = R$ to ensure there are no logs in the anomalous dimension. For C_0 ,

$$R \frac{d}{dR} \ln C_0(Q, R, R) = \bar{\gamma}[\alpha_s(R)] - \left(\frac{R}{Q} \right)^p \gamma[\alpha_s(R)], \quad (10)$$

where $\bar{\gamma}[\alpha_s] = \sum_{n=0}^{\infty} \bar{\gamma}_n[\alpha_s(R)/4\pi]^{n+1}$ and $\gamma[\alpha_s] = \sum_{n=0}^{\infty} \gamma_n[\alpha_s(R)/4\pi]^{n+1}$ are the $\overline{\text{MS}}$ and R anomalous dimensions. Here $\gamma_{n-1} = p a_{n0} - 2 \sum_{m=1}^{n-1} m a_{m0} \beta_{n-m-1}$ and we are using the $\overline{\text{MS}}$ β -function $\mu d/d\mu \alpha_s(\mu) = -\alpha_s^2(\mu)/(2\pi) \sum_{n=0}^{\infty} \beta_n[\alpha_s(\mu)/4\pi]^n$. The choice in Eq. (8) keeps Eq. (10) simple. In cases where $\bar{\gamma}$ is absent we expect Eq. (10) to converge to lower scales due to the $(R/Q)^p$ factor multiplying γ . For $R_1 > R_0$ the solution of Eq. (10) is $[U_\mu = U_\mu(R_1, R_0)]$

$$C_0(Q, R_0, R_0) = C_0(Q, R_1, R_1) U_R(Q, R_1, R_0) U_\mu, \quad (11)$$

where U_μ is a usual $\overline{\text{MS}}$ evolution factor and U_R is the R -evolution. For $p = 1$ the complete solution for U_R was obtained in Ref. [6]. It is straightforward to generalize this to any p . At $N^{k+1}\text{LL}$ order the (real) result is

$$U_R(Q, R_1, R_0) = \exp \left\{ \left(\frac{\Lambda_{\text{QCD}}^{(k)}}{Q} \right)^p \sum_{j=0}^k S_j (-p)^j e^{i\pi p \hat{b}_1} \times p^{(p\hat{b}_1)} [\Gamma(-p\hat{b}_1 - j, pt_0) - \Gamma(-p\hat{b}_1 - j, pt_1)] \right\}, \quad (12)$$

with $\Gamma(c, t)$ the incomplete gamma function and $t_{0,1} = -2\pi/(\beta_0 \alpha_s(R_{0,1}))$. $\Lambda_{\text{QCD}}^{(0)} = Re^t$, $\Lambda_{\text{QCD}}^{(1)} = Re^t(-t)^{\hat{b}_1}$, and $\Lambda_{\text{QCD}}^{(2)} = Re^t(-t)^{\hat{b}_1} e^{-\hat{b}_2/t}$ are evaluated at a large reference R with $t = -2\pi/(\beta_0 \alpha_s(R))$, and $\hat{b}_1 = \beta_1/(2\beta_0^2)$, $\hat{b}_2 = (\beta_1^2 - \beta_0 \beta_2)/(4\beta_0^4)$, and $\hat{b}_3 = (\beta_1^3 - 2\beta_0 \beta_1 \beta_2 + \beta_0^2 \beta_3)/(8\beta_0^6)$. Defining $\tilde{\gamma}_n = \gamma_n/(2\beta_0)^{n+1}$ the coefficients of U_R needed for the first three orders of R -evolution are

$$S_0 = \tilde{\gamma}_0, \quad S_1 = \tilde{\gamma}_1 - (\hat{b}_1 + p\hat{b}_2) \tilde{\gamma}_0, \quad (13)$$

$$S_2 = \tilde{\gamma}_2 - (\hat{b}_1 + p\hat{b}_2) \tilde{\gamma}_1 + [(1 + p\hat{b}_1) \hat{b}_2 + p(p\hat{b}_2^2 + \hat{b}_3)/2] \tilde{\gamma}_0.$$

Eq. (6) becomes $C_0(Q, R_1, R_1) U_R(Q, R_1, R_0) U_\mu(R_1, R_0) \times \theta_0(R_0) + \theta_1(R_0, R_0)/Q^p$, and this result sums logs between $R_1 \sim Q$ and $R_0 \sim \Lambda_{\text{QCD}}$. This gives natural R scales for coefficients and matrix elements in the OPE.

Heavy Meson Mass Splittings in MSR

The $\overline{\text{MS}}$ OPE for the mass-splitting of heavy mesons, $\Delta m_H^2 = m_{H^*}^2 - m_H^2$ for $H = B, D$, is $\Delta m_H^2 = \bar{C}_G(m_Q, \mu) \mu_G^2(\mu) + \sum_i \bar{C}_i(m_Q, \mu) 2\rho_i^3(\mu)/(3m_Q) + \mathcal{O}(\Lambda_{\text{QCD}}^3/m_Q^2)$, where $m_Q = m_b$ or m_c . Here $\mu_G^2 = -\langle B_v | \bar{h}_v g \sigma_{\mu\nu} G^{\mu\nu} h_v | B_v \rangle / 3$ is the matrix element of the chromomagnetic operator, and ρ_i^3 for $i = \pi G, A, LS, \bar{\Lambda} G$ are $\mathcal{O}(\Lambda_{\text{QCD}}^3)$ matrix elements [7], with $\rho_{\bar{\Lambda} G}^3(\mu) = (3/2)\bar{\Lambda}\mu_G^2(\mu)$. At the order of our analysis tree level values for the $\bar{C}_i$ suffice, so with $\bar{\Sigma}_\rho(\mu) = (2/3)[\rho_{\pi G}^3(\mu) + \rho_A^3(\mu) - \rho_{LS}^3(\mu) + \rho_{\bar{\Lambda} G}^3(\mu)]$,

$$\Delta m_H^2 = \bar{C}_G(m_Q, \mu) \mu_G^2(\mu) + \bar{\Sigma}_\rho(\mu)/m_Q + \dots \quad (14)$$

Taking the ratio of mass splittings $r = \Delta m_B^2/\Delta m_D^2$ gives

$$r = \frac{\bar{C}_G(m_b, \mu)}{\bar{C}_G(m_c, \mu)} + \frac{\bar{\Sigma}_\rho(\mu)}{\mu_G^2(\mu)} \left(\frac{1}{m_b} - \frac{1}{m_c} \right) + \dots \quad (15)$$

The first term in this OPE gives a purely perturbative prediction for r . $\bar{C}_G$ is known to suffer from an $\mathcal{O}(\Lambda_{\text{QCD}}/m_Q)$ infrared renormalon ambiguity [7], with a corresponding ambiguity in $\bar{\Sigma}_\rho(\mu)$. The three-loop computation of Ref. [8] yields, $r = 1 - 0.1113|_{\alpha_s} - 0.0780|_{\alpha_s^2} - 0.0755|_{\alpha_s^3}$ at fixed order with $\mu = m_c$, and $r = (0.8517)_{\text{LL}} + (-0.0696)_{\Delta\text{NLL}} + (-0.0908)_{\Delta\text{NNLL}}$ in RGE-improved perturbation theory, with no sign of convergence in either case. In $\overline{\text{MS}}$ these leading predictions are unstable due to the $p = 1$ renormalon in $\bar{C}_G$.

Lets examine the analogous result in the MSR scheme

$$\Delta m_H^2 = C_G(m_Q, R, \mu) \mu_G^2(\mu) + \frac{\Sigma_\rho(R, \mu)}{m_Q} + \dots \quad (16)$$

Since $p = 1$ the MSR definition in Eq. (9) gives

$$C_G(m_Q, R, \mu) \equiv \bar{C}_G(m_Q, \mu) [\bar{C}_G(R, \mu)]^{-R/m_Q}, \quad (17)$$

where $\bar{C}_G(m, \mu)$ is obtained from Ref. [8] and we expand in $\alpha_s(\mu)$. The OPE in MSR at a scale $R_0 \gtrsim \Lambda_{\text{QCD}}$ gives

$$r = \frac{C_G(m_b, R_0, R_0)}{C_G(m_c, R_0, R_0)} + \frac{\Sigma_\rho(R_0, R_0)}{\mu_G^2(R_0)} \left(\frac{1}{m_b} - \frac{1}{m_c} \right). \quad (18)$$

Large logs in $C_G(m_Q, R_0, R_0)$ can be summed with the R-RGE in Eqs. (11–13). For simplicity we integrate out the b and c -quarks simultaneously at a scale $R_1 \simeq \sqrt{m_b m_c} \gg R_0 \simeq \Lambda_{\text{QCD}}$. This scale for R_1 keeps $\ln(R_1/m_{b,c})$ small. With R-evolution and U_R from Eq. (12) we have

$$r = \frac{C_G(m_b, R_1, R_1) U_R(m_b, R_1, R_0)}{C_G(m_c, R_1, R_1) U_R(m_c, R_1, R_0)} + \frac{\Sigma_\rho(R_0, R_0)}{\mu_G^2(R_0)} \left(\frac{1}{m_b} - \frac{1}{m_c} \right). \quad (19)$$

This expression is independent of R_1 and R_0 . Order-by-order, varying R_1 about $\sqrt{m_b m_c}$ yields an estimate of higher order perturbative uncertainties, much like varying μ in $\overline{\text{MS}}$. For R_0 the dependence cancels between the




FIG. 1: Perturbative predictions at leading order in $1/m_Q$ for the ratio r of the B - B^* and D - D^* mass splittings in the MSR-scheme (solid) versus $\overline{\text{MS}}$ (dashed). The R_0 dependence of the solid red curve provides an estimate for the power correction, independent of the comparison with the experimental data. Neither R_1 nor μ variation is shown in the figure.

first term in r and the Σ_ρ power correction. In MSR the term $\Sigma_\rho(R_0, R_0)$ is $\sim \Lambda_{\text{QCD}}^3$ and can be positive or negative. One may expect that there is a value of R_0 where $\Sigma_\rho(R_0, R_0)$ vanishes. Thus keeping only the first term in Eq. (19) and varying $R_0 \gtrsim \Lambda_{\text{QCD}}$ yields an estimate for the size of this power correction. This technique goes beyond the dimensional analysis estimates used in $\overline{\text{MS}}$.

Fig. 1 gives perturbative predictions for r at different orders using the first terms in Eqs. (15,19) with $m_b = 4.7$ GeV, $m_c = 1.6$ GeV, $\alpha_s(\sqrt{m_b m_c}) = 0.2627$ and the 4-loop β -function. The solid lines are from the MSR scheme, plotted as functions of R_0 . The dashed lines are the fixed order $\overline{\text{MS}}$ results with $\mu = \sqrt{m_b m_c}$. The MSR results exhibit a dramatic improvement in convergence over $\overline{\text{MS}}$ for a wide range of R_0 values. Varying $R_1 = \sqrt{m_b m_c}/2$ to $2\sqrt{m_b m_c}$ at $N^3\text{LL}(\text{MSR})$ gives $\Delta r \simeq \pm 0.008$, which is a significant improvement over μ variation in the same range for $N^3\text{LO}(\overline{\text{MS}})$ where $\Delta r \simeq \pm 0.068$. The MSR results converge to an R_0 dependent curve, whose dependence cancels against $\Sigma_\rho(R_0, R_0)$, so the residual R_0 dependence provides a method to estimate the size of this power correction. The range $R_0 = 0.7$ GeV to 1.2 GeV keeps R_0 below m_c and above Λ_{QCD} and yields

$$r = 0.860 \pm (0.065)_{\Sigma_\rho} \pm (0.008)_{\text{pert.}} \quad (20)$$

This estimate for the Σ_ρ power correction in MSR is in good agreement with experiment, $r_{\text{expt}} = 0.886$ ($D_{u,d}^{(*)}$, $B_{u,d}^{(*)}$) and 0.854 ($D_s^{(*)}$, $B_s^{(*)}$). MSR achieves a convergent perturbative prediction for r at leading order in the OPE, and a $1/m_Q$ power correction of moderate size, ~ 0.065 , significantly smaller than the dimensional analysis estimate of $\Lambda_{\text{QCD}}(1/m_c - 1/m_b) \sim 0.2$ in $\overline{\text{MS}}$.

Ellis-Jaffe sum rule in MSR

In $\overline{\text{MS}}$ the Ellis-Jaffe sum rule [9] for the proton in DIS with momentum transfer Q is $M_1(Q) = [\bar{C}_B(Q, \mu) \theta_B + \bar{C}_0(Q, \mu) \hat{a}_0/9] + \theta_1(\mu)/Q^2$. $\bar{C}_{B,0}$ are




FIG. 2: Perturbative results for the Ellis-Jaffe sum rule in the MSR, RS, and $\overline{\text{MS}}$ schemes, at leading order in $1/Q$. For all curves the one parameter, $\hat{a}_0$, is fixed by data at $Q \simeq 5$ GeV.




FIG. 3: Uncertainty estimates in the MSR scheme and $\overline{\text{MS}}$ scheme for the Ellis-Jaffe sum rule at leading order in $1/Q$.

known at 3 loops [10]. The two leading order terms are written so that both coefficients and matrix elements are separately μ -independent: $\theta_B = g_A/12 + a_8/36$ is given by the axial couplings $g_A = 1.2694$ and $a_8 = 0.572$ for the nucleon and hyperon, while $\hat{a}_0$ is a Q independent $\overline{\text{MS}}$ matrix element. θ_1 denotes all $1/Q^2$ power corrections with their Wilson coefficients at tree level. The $\overline{\text{MS}}$ coefficients are affected by a $p = 2$ renormalon [11], which is removed in the MSR scheme. Eq. (9) gives $[i = B, 0]$

$$C_i(Q, R, R) \equiv \bar{C}_i(Q, R) [\bar{C}_i(R, R)]^{-R^2/Q^2}. \quad (21)$$

With R -evolution the MSR OPE prediction is

$$M_1(Q) = [C_B(Q, R_1, R_1)U_R^B(Q, R_1, R_0)\theta_B + C_0(Q, R_1, R_1)U_R^0(Q, R_1, R_0)\hat{a}_0/9] + \theta_1(R_0, R_0)/Q^2, \quad (22)$$

where $U_R^{B,0}$ are given by Eq. (12) with $p = 2$ and the corresponding $(a_{n0})^{B,0}$ determine the appropriate $(\gamma_n)^{B,0}$.

Figures 2,3 show perturbative predictions for the Ellis-Jaffe sum rule at leading power in $1/Q$, compared with proton data from Ref. [12]. We use $\alpha_s(4 \text{ GeV}) = 0.2282$, and the 4-loop β with 4 flavors. In Fig. 2, we show order-by-order results for the $\overline{\text{MS}}$ scheme at $\mu = Q$, and for the resummed MSR scheme with $R_1 = Q$ and $R_0 = 0.9$ GeV. We fix $\hat{a}_0 = 0.141$ so that $\overline{\text{MS}}$ and MSR agree with the data for $Q \simeq 5$ GeV. $\overline{\text{MS}}$ agrees well with the data for large Q , but turns away at $Q \lesssim 2$ GeV and no longer converges. In contrast the MSR results still converge quickly and exhibit excellent agreement with the data over a wide range of Q values. The NLL MSR result already has the right curvature, and the NNLL and $N^3\text{LL}$ curves further improve the agreement. We also include predictions in the RS scheme with subtraction scale $\nu_f = 1.0$ GeV from Fig.3d of Ref. [5], which improve slightly over the $\overline{\text{MS}}$ results, but may not be capturing the dominant power law dependence on the factorization scale. In Fig. 3 we show uncertainties for three loop results in the $\overline{\text{MS}}$ and MSR schemes. The dashed red curve is the $\overline{\text{MS}}$ prediction, and the blue band estimates the higher-order perturbative uncertainties varying μ in the range $\mu^{\min}(Q) < \mu < 2Q$. For $Q > 1.5$ GeV, $\mu^{\min} = Q/2$, while for $Q < 1.5$ GeV, $\mu^{\min} = 1.3Q/(1.1 + Q/(1 \text{ GeV}))$. The red solid line is the MSR prediction, the red band is the perturbative uncertainty from varying R_1 in the same range as was done for μ in $\overline{\text{MS}}$, and the green band estimates the $1/Q^2$ power correction by varying $R_0 = 0.7$ to 1.2 GeV. Fig. 3 implies $-0.01 \text{ GeV}^2 \lesssim \theta_1(R_0, R_0) \lesssim 0.01 \text{ GeV}^2$ in MSR, which is a much smaller power correction than the $\sim 0.1 \text{ GeV}^2$ estimate obtained from naive dimensional analysis in $\overline{\text{MS}}$.

This work was supported by the EU network, MRTN-CT-2006-035482 (Flavianet), Spanish Ministry of Education, FPA2008-00592, the Office of Nuclear Physics of the U.S. Department of Energy, DE-FG02-94ER40818, the Alexander von Humboldt foundation, and the Max-Planck-Institut für Physik guest program.

- [1] M. Beneke, Phys. Rept. **317**, 1 (1999).
- [2] A. H. Mueller, Nucl. Phys. **B250**, 327 (1985).
- [3] M. Luke et al., Phys. Rev. **D51**, 4924 (1995).
- [4] L. Maiani et al., Nucl. Phys. **B368**, 281 (1992).
- [5] F. Campanario and A. Pineda, Phys. Rev. **D72**, 056008 (2005); A. Pineda, JHEP **06**, 022 (2001).
- [6] A. H. Hoang, A. Jain, I. Scimemi, and I. W. Stewart, Phys. Rev. Lett. **101**, 151602 (2008), 0803.4214.
- [7] A. Grozin and M. Neubert, Nucl. Phys. **B508**, 311 (1997).

- [8] A. G. Grozin et al., Nucl. Phys. **B789**, 277 (2008).
- [9] J. R. Ellis and R. L. Jaffe, Phys. Rev. **D9**, 1444 (1974).
- [10] S. A. Larin, T. van Ritbergen, and J. A. M. Vermaseren, Phys. Lett. **B404**, 153 (1997).
- [11] D. J. Broadhurst and A. L. Kataev, Phys. Lett. **B315**, 179 (1993).
- [12] M. Osipenko et al., Phys. Rev. **D71**, 054007 (2005).